

A Reliable Research Partner in Life Science and Medicine

APC Anti-Human CD33 Antibody[P67.6]

Catalog Number: AN00333E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_	e				40		
		•	^		TI.	$\boldsymbol{\cap}$	
	4-	-	•	w	ш		

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ

Clone No. P67.6

Isotype Control APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]

Conjugation APC

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with APC Anti-Human CD33 Antibody[P67.6] (filled gray histogram) or APC Mouse IgG1, κ Isotype Control (empty black histogram). Cells in the monocytes gate were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

Antigen Information

Alternate Names Myeloid cell surface antigen CD33;CD33;Sialic acid-binding lg-like lectin 3;Siglec-3;

Web: www.elabscience.cn

gp67;SIGLEC3

 Uniprot ID
 P20138

 Gene ID
 945

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD33 is a 67 kD type I transmembrane glycoprotein also known as Siglec-3, gp67, and p67. It is a sialoadhesion immunoglobulin superfamily member expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. CD33 is absent on normal platelets, lymphocytes, erythrocytes and hematopoietic stem cells. CD33 functions as a sialic acid-dependent cell adhesion molecule with carbohydrate/lectin binding activity.

Rev. V1.5

Web: www.elabscience.cn