

A Reliable Research Partner in Life Science and Medicine

# CD27/TNFRSF7 Polyclonal Antibody(Capture/Detector)

catalog number: AN004010P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$		cription		
	OC.	cri		OT:
$\mathbf{L}$	CO.		JU	ш

Reactivity Human; Mouse; Rat

Immunogen Recombinant Mouse CD27/TNFRSF7 Protein expressed by Mammalian

Host Rabbit
Isotype Rabbit IgG

**Purification** Antigen Affinity Purification

**Conjugation** Unconjugated


**Buffer** Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

## Applications Recommended Dilution

 ELISA Capture
 2-8 μg/mL

 ELISA Detector
 0.1-0.4 μg/mL

#### Data



Sandwich ELISA-Recombinant Mouse CD27/TNFRSF7
Protein standard curve.Background subtracted standard curve using Anti-CD27/TNFRSF7 antibody(AN004010P)
(Capture),Anti-CD27/TNFRSF7 antibody(AN004010P)
(Detector).The reference range value is 62.5 - 4,000 pg/mL for mouse.

## **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

### Background

CD27, also known as TNFRSF7, is a member of the TNF-receptor superfamily limited to cells of the lymphoid lineage, and exists as both a dimeric glycoprotein on the cell surface and as a soluble protein in serum. As a type I transmembrane glycoprotein of about 55 kDa existing as disulfide-linked homodimer, CD27 has been shown to play roles in lymphoid proliferation, differentiation, and apoptosis. It has an important role in the generation of T cell immunity and is an robust marker for normal memory B cells. It is a T and B cell co-stimulatory molecule, the activity of CD27 is governed by its TNF-like ligand CD70 on lymphocytes and dendritic cells. The CD27-CD70 interaction is required for Th1 generation responses to differentiation signals and long-term maintenance of T cell immunity, and meanwhile, plays a key role in regulating B-cell differentiation, activation and immunoglobulin synthesis.

#### For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email:techsupport@elabscience.cn