# Purified Anti-Human CD45 Antibody[BC8]

**catalog number:** **E-AB-F13720P**


**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Reactivity** | Human |
| **Immunogen** | Recombinant Human CD45 protein |
| **Host** | Mouse |
| **Isotype** | Mouse IgG1, κ |
| **Clone** | BC8 |
| **Purification** | >98%, Protein A/G purified |
| **Buffer** | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

| Applications | Recommended Dilution |
|---|---|
| **FCM** | ≤ 0.2 μg per million cells in 100 μL volume |

## Data



Human peripheral blood were stained with 0.2 μg Purified Anti-Human CD45 Antibody[BC8] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

## Preparation & Storage

| | |
|---|---|
| **Storage** | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| **Shipping** | Ice bag |

## Background

CD45, previously called LCA (leukocyte common antigen), T200, or Ly5 in mice, is member C of the class 1 (receptor-like) protein tyrosine phosphatase family (PTPRC). It is a variably glycosylated 180-220 kDa transmembrane protein that is abundantly expressed on all nucleated cells of hematopoietic origin . Multiple splicing isoforms of exon 4 (A), 5 (B), and 6 (C) are expressed according to cell type, developmental stage and antigenic exposure . The longest form, CD45RABC (called B220 in mouse) is expressed on B lymphocytes, The shortest form, CD45R0, lacking exons 4, 5 and 6 which encode aa 34-194, is expressed on memory cells, while intermediate sizes are expressed on other T


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com