

AMPK alpha1 Monoclonal Antibody

catalog number: E-AB-22125

Note: Centrifuge before opening to ensure complete recovery of vial contents.



Description

| | |
|---------------------|---|
| Reactivity | Human;Mouse;Rat |
| Immunogen | Synthetic Peptide |
| Host | Mouse |
| Isotype | IgG |
| Clone | 6B6 |
| Purification | Protein A purification |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol. |

Applications

| Applications | Recommended Dilution |
|--------------|----------------------|
| WB | 1:1000-2000 |
| IHC | 1:50-100 |

Data



Western Blot analysis of 1) Hela, 2) 293T, 3) 3T3, 4) PC-12 cells using AMPK alpha1 Monoclonal Antibody at dilution of 1:2000. Immunohistochemistry of paraffin-embedded Mouse colon tissue using AMPK alpha1 Monoclonal Antibody at dilution of 1:200.

Observed-MW:63 kDa

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

Responsible for the regulation of fatty acid synthesis by phosphorylation of acetyl-CoA carboxylase. It also regulates cholesterol synthesis via phosphorylation and inactivation of hormone-sensitive lipase and hydroxymethylglutaryl-CoA reductase. Appears to act as a metabolic stress-sensing protein kinase switching off biosynthetic pathways when cellular ATP levels are depleted and when 5'-AMP rises in response to fuel limitation and/or hypoxia. This is a catalytic subunit.

For Research Use Only