

Recombinant Human NGAL/Lipocalin-2 Protein (His Tag, Human Cells)

Catalog Number: PKSH032806

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human NGAL;Lipocalin-2 protein Gln21-Gly198, with an C-terminal His
Calculated MW	21.6 kDa
Observed MW	23 kDa
Accession	P80188
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/ gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 µm filtered solution of PBS, 50% Glycerol, pH 7.4.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

LCN2 is iron-trafficking protein involved in multiple processes such as apoptosis; innate immunity and renal development. LCN2 binds iron through association with 2;5-dihydroxybenzoic acid (2;5-DHBA); a siderophore that shares structural similarities with bacterial enterobactin; and delivers or removes iron from the cell; depending on the context. LCN2 is involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular iron concentration without promoting apoptosis; while iron-free form decreases intracellular iron levels; inducing expression of the proapoptotic protein BCL2L1/BIM; resulting in apoptosis. LCN2 is involved in innate immunity; possibly by sequestering iron; leading to limit bacterial growth.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017