

## Recombinant Human BCMA/TNFRSF17 Protein (Fc Tag)

**Catalog Number:** PKSH033752

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human BCMA;TNFRSF17 protein Met1-Ala54, with an C-terminal Fc
<b>Mol_Mass</b>	32.8 kDa
<b>Accession</b>	Q02223
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

B cell maturation protein (BCMA) is a type III membrane protein which belongs to the TNF receptor superfamily (TNFRSF17). BCMA contains one extracellular cysteine rich domain; as well as TACI. Human BCMA is a 184 amino acid (aa) protein consisting of a 54 aa extracellular domain; a 23 aa transmembrane domain; and a 107 aa intracellular domain. Mouse and human BCMA share 62% amino acid identity. BCMA is mainly expressed in immune organs and mature B cell lines. BCMA appears to be localized to the Golgi compartment. BCMA may play an important role in B cell development; function and regulation.

### For Research Use Only