

Purified Anti-Human/Mouse CD44 Antibody[IM7], Functional Grade

catalog number: E-AB-F11000

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse
Immunogen	Recombinant Human;Mouse CD44 protein
Host	Rat
Isotype	Rat IgG2b, κ
Clone	IM7
Purification	>98%, Protein A/G purified
Buffer	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

Applications


Recommended Dilution

FCM	$\leq 0.2 \mu\text{g}$ per million cells in 100 μL volume
Neut	Reported in the literature

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μg Purified Anti-Human/Mouse CD44 Antibody[IM7], Functional Grade (Right) and 0.2 μg Rat IgG2b, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.



C57/BL6 Mouse splenocytes were stained with 0.2 μg Purified Anti-Human/Mouse CD44 Antibody[IM7], Functional Grade (Right) and 0.2 μg Rat IgG2b, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

For Research Use Only

CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from low or mid levels to high levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. High CD44 expression on Treg cells has been associated with potent suppressive function via high production of IL-10. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

Application References

Luca G Guidotti, et al. Cell. 2015 Apr 23;161(3):486-500.