

Recombinant Human TPBG/5T4 (C-mFc)

Catalog Number: PKSH033920

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human TPBG/5T4 protein Ser32-Ser355, with an C-terminal mFc
Calculated MW	61.7 kDa
Observed MW	80-110 kDa
Accession	Q13641
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Trophoblast glycoprotein, also known as TPBG and 5T4, it belongs to the LRR family of proteins. The C-terminus of LRR N-terminal cap (LRRNT) and LRR 1 are essential for the inhibition of the Wnt signaling pathway. TPBG may function as an inhibitor of Wnt/beta-catenin signaling by indirectly interacting with LRP6 and blocking Wnt3a-dependent LRP6 internalization. It expressed by all types of trophoblasts as early as 9 weeks of development. In adult tissues, the expression is limited to a few epithelial cell types but is found on a variety of carcinoma.

For Research Use Only