# Recombinant Human CD27 Ligand/TNFSF7/CD70 (C-mFc)

Catalog Number: PKSH033858



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Des	cription

 Species
 Human

 Mol\_Mass
 43.0 kDa

 Accession
 P32970

**Bio-activity** Not validated for activity

# **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

CD27 Ligand (CD70) is a 30 kDa type 2 transmembrane glycoproteinis that belongs to the tumor necrosis factor (TNF) ligand family. Human CD27 Ligand extracellular region is 64% as identical to mouse and rat CD27 Ligand extracellular regions. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis.

## For Research Use Only