## Recombinant Human Mesothelin/MSLN Protein (aa 37-286, Fc Tag)

Catalog Number: PKSH032747



Description			
Species	Human		
Mol_Mass	53.9 kDa		
Accession	Q13421		
<b>Bio-activity</b>	Not validated for activity		
Properties			
Purity	> 90 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		

Note: Centrifuge before opening to ensure complete recovery of vial contents.

kDa	MK	ence R	
120 90			
60		-	e°.
40			
30	-		ence
20			
14			

> 90 % as determined by reducing SDS-PAGE.

## Background

Data

Mesothelin is a cell surface glycoprotein whose expression is limited to mesothelial cells of the serosa (pleura; pericardium; and peritoneum) and epithelial cells of the trachea; tonsils; fallopian tube; and kidneys. Mesothelin plays an important role in cell survival; proliferation; migration; invasion; tumor progression; and resistance to chemotherapy. The overexpression of mesothelin can activate NF- $\kappa$ B and signal transducer and activator of transcription 3 (Stat3); inhibit apoptotic signaling and TNF- $\alpha$ -induced apoptosis; and accelerate the Gl–S transition. Mesothelin is also found overexpressed in various cancers; including malignant mesothelioma; pancreatic or ovarian carcinoma; sarcomas and in some gastrointestinal or pulmonary carcinomas. As a result of its limited expression in normal tissues; mesothelin has been reported as an ideal tumor-associated marker for the development of targeted therapy.

## For Research Use Only