

Recombinant Human Biglycan/BGN protein (His Tag)

Catalog Number: PDMH100070

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Biglycan;BGN protein Glu20-Lys368, with an C-terminal His
Calculated MW	40.4 kDa
Observed MW	42 kDa
Accession	P21810
Bio-activity	Not validated for activity

Properties

Purity	> 80% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human Biglycan/BGN proteins,
2µg/lane of Recombinant Human Biglycan/BGN proteins
was resolved with SDS-PAGE under reducing conditions,
showing bands at 42 KD.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V4.0

Biglycan, also known as PG-S1 and BGN, is a small leucine-rich repeat proteoglycan (SLRP). It can be detected in a variety of extracellular matrix tissues, including bone, cartilage and tendon. Biglycan consists of a protein core containing leucine-rich repeat regions and two glycosaminoglycan (GAG) chains consisting of either chondroitin sulfate (CS) or dermatan sulfate (DS). Non-glycanated forms of biglycan (no GAG chains) increase with age in human articular cartilage. Biglycan interacts with collagen, both via the core protein and GAG chains. Biglycan plays a role in the mineralisation of bone. Biglycan core protein binds to the growth factors BMP-4 and influences its bioactivity.