

A Reliable Research Partner in Life Science and Medicine

Recombinant Human Fcy RIIB/CD32b Protein(His Tag)

Catalog Number: PDMH100315

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human Fcy RIIB/CD32b proteins Ala46-Pro217, with an C-

terminal His

Calculated MW 18.8 kDa
Observed MW 30 kDa
Accession P31994

Bio-activity Not validated for activity

Properties

Purity > 90% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping

This product is provided as lyophilized powder which is shipped with ice packs.

Formulation


Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human Fc γ RIIB/CD32b proteins , 2 μ g/lane of Recombinant Human Fc γ RIIB/CD32b proteins was resolved with SDS-PAGE under reducing conditions , showing bands at 30 KD

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

FcyRIIB is a low affinity receptor that recognizes the Fc portion of IgG. The human CD32 group consists of FcyRIIA, Fcy RIIB, and FcyRIIC proteins that share 94-99% sequence identity in their extracellular domains but differ substantially in their transmembrane and cytoplasmic domains. FcyRII protein is expressed on cells of both myeloid and lymphoid lineages as well as on cells of non-hematopoietic origin. FcyRIIB has an intrinsic cytoplasmic immunoreceptor tyrosine-based inhibitory motif (ITIM) and delivers an inhibitory signal upon ligand binding. Ligation of FcyRIIB on B cells down-regulates antibody production and in some circumstances may promote apoptosis. Co-ligation of FcyRIIB on dendritic cells inhibits maturation and blocks cell activation. FcyRIIB may also be a target for monoclonal antibody therapy for malignancies. FcyRIIB plays an important negative-regulating role through modulating the signals from activation receptors.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com