Elabscience®

VTI1A Polyclonal Antibody


catalog number: E-AB-63813

Note: Centrifuge before opening to ensure complete recovery of vial contents.

1:500-1:2000

| Applications | Recommended Dilution |
|--------------|--|
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Purification | Affinity purification |
| Isotype | IgG |
| Host | Rabbit |
| Immunogen | Recombinant fusion protein of human VTI1A (NP_660207.2). |
| Reactivity | Human;Mouse;Rat |
| Description | |

WB Data



Western blot analysis of extracts of various cell lines using

VTI1A Polyclonal Antibody at dilution of 1:1000.

Observed-MW:30 kDa Calculated-MW:23 kDa/25 kDa

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |

Background

The protein encoded by this gene is a member of the family of soluble N-ethylmaleimide-sensitive fusion proteinattachment protein receptors (SNAREs) that function in intracellular trafficking. This family member is involved in vesicular transport between endosomes and the trans-Golgi network. It is a vesicle-associated SNARE (v-SNARE) that interacts with target membrane SNAREs (t-SNAREs). Polymorphisms in this gene have been associated with binocular function, and also with susceptibility to colorectal and lung cancers. A recurrent rearrangement has been found between this gene and the transcription factor 7-like 2 (TCF7L2) gene in colorectal cancers. Alternative splicing results in multiple transcript variants.

For Research Use Only