

TARC/CCL17 (C-6His), Human, Recombinant**Cat. No. : GPCK109****产品信息**

物种	Human
表达宿主	Human Cells
序列信息	Ala24-Ser94
检索号	Q92583
分子量	9.1 kDa
有效期	12 months

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl, pH 8.0.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

C-C Motif Chemokine 17 (CCL17) is a novel CC Chemokine, it belongs to the intercrine beta (Chemokine CC) family. CCL17 is expressed at high levels in thymus, and at a lower level in lung, colon, and small intestine. CCL17 is also transiently expressed in stimulated peripheral blood mononuclear cells. Among CC Chemokine family members, CCL17 has approximately 24-29% amino acid sequence identity with RANTES, MIP-1 alpha, MIP-1 beta, MCP-1, MCP-2 and MCP-3. CCL17 has been identified to be Chemotactic factor for T-lymphocytes but not monocytes or granulocytes. CCL17 plays a role in T-cell development in thymus and in trafficking and activation of mature T-cells.

