# **Elabscience**®

# **IRGC Polyclonal Antibody**

## catalog number: E-AB-18127


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Rat
Immunogen	Synthetic peptide of human IRGC
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution

 Appreadors
 Recommended Dirution

 IHC
 1:30-1:150

### Data



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using IRGC Polyclonal Antibody at dilution of 1:30(×200) Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using IRGC Polyclonal Antibody at dilution of 1:30(×200)

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the
	temperature recommended.

### Background

Immunity-related GTPases (IRG) (also known as p47 GTPases) are a family of GTPase proteins found in vertebrates, which play critical roles in mediating innate resistance to intracellular pathogens. IRG genes have been found in a number of mammals and lower species including mice, rats, zebrafish and humans. Most of the mouse genes contain interferon-stimulated response elements which mediate transcriptional activation by IFNs. In humans, only two IRG genes have been found: human IRGC encodes a full-length IRG protein that, like the mouse homologue, is constitutively expressed in testis, while human IRGM encodes a considerably truncated protein that is constitutively expressed in cultured cells including some macrophage cell lines. As the two human genes IRGC and IRGM are not subject to IFN control, it has been suggested that the host resistance mechanism supported by IRG proteins in the mouse is lacking in humans.