Recombinant Mouse IL22 protein (His Tag)

Catalog Number: PDEM100268



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 10	00	cri	m	17	٦m
u		\mathbf{u}		7 U	

Species Mouse

Source E.coli-derived Mouse IL22 protein Leu34-Val179, with an N-terminal His

 Mol_Mass
 16.0 kDa

 Accession
 Q9JJY9

Bio-activity Not validated for activity

Properties

Purity > 95% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-22 (IL-22) was initially identified as a gene induced by IL-9 in mouse T cells and mast cells. Mouse IL-22 cDNA encodes a 179 amino acid residue protein with a putative 33 amino acid signal peptide that is cleaved to generate a 147 amino acid mature protein that shares approximately 79% and 22% sequence identity with human IL22 and IL10, respectively. IL22 has been shown to activate STAT-1 and STAT-3 in several hepatoma cell lines and up-regulate the production of acute phase proteins. IL-22 is produced by normal mouse T cells upon Con A activation. Mouse IL-22 expression is also induced in various organs upon lipopolysaccharide injection, suggesting that IL-22 may be involved in inflammatory responses. The functional IL-22 receptor complex consists of two receptor subunits, IL-22R (previously an orphan receptor named CRF2-9) and IL-10R β (previously known as CRF2-4), belonging to the class II cytokine receptor family.

For Research Use Only