

Recombinant Human PHPT1/PHP14 Protein (His Tag)

Catalog Number: PKSH030832

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human PHPT1/PHP14 protein Ala 2-Tyr 125, with an N-terminal His
Calculated MW	15.2 kDa
Observed MW	15.2 kDa
Accession	Q9NRX4-1
Bio-activity	Not validated for activity

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

PHPT1, also known as 14 kDa phosphohistidine phosphatase, phosphohistidine phosphatase 1, protein janus-A homolog, PHP14, is a cytoplasm protein which belongs to the janus family. PHPT1 / PHP14 is expressed abundantly in heart and skeletal muscle. Phosphatases are a diverse group of enzymes that regulate numerous cellular processes. Much of what is known relates to the tyrosine, threonine, and serine phosphatases, whereas the histidine phosphatases have not been studied as much. Protein histidine phosphorylation exists widely in vertebrates, and it plays important roles in signal transduction and other cellular functions. Protein histidine phosphorylation accounts for about 6% of the total protein phosphorylation in eukaryotic cells. The knowledge about eukaryotic PHPT (protein histidine phosphatase) is still very limited. To date, only one vertebrate PHPT has been discovered, and two crystal structures of human PHPT1 have been solved. PHPT1 / PHP14 can dephosphorylate a variety of proteins (e.g. ATP-citrate lyase and the beta-subunit of G proteins). A putative active site has been identified by its electrostatic character, ion binding, and conserved protein residues.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017