


## CD73/NT5E Polyclonal Antibody

catalog number: E-AB-93185

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |                                                                                    |
|--------------|------------------------------------------------------------------------------------|
| Reactivity   | Human;Mouse;Rat                                                                    |
| Immunogen    | Recombinant fusion protein of human CD73/NT5E                                      |
| Host         | Rabbit                                                                             |
| Isotype      | Rabbit IgG                                                                         |
| Purification | Affinity purification                                                              |
| Buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution                                                               |
| WB           | 1:500-1:2000                                                                       |
| IHC          | 1:50-1:200                                                                         |


## Data




Western blot analysis of extracts of various cell lines using CD73/NT5E Polyclonal Antibody at 1:1000 dilution.

**Observed-MW:70 kDa**


**Calculated-MW:57 kDa/63 kDa**



Immunohistochemistry of paraffin-embedded rat kidney using CD73/NT5E Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded human placenta using CD73/NT5E Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded mouse testis using CD73/NT5E Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

## Preparation & Storage

|                 |                                                                                                          |
|-----------------|----------------------------------------------------------------------------------------------------------|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.                                          |
| <b>Shipping</b> | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

## Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.8

The protein encoded by this gene is a plasma membrane protein that catalyzes the conversion of extracellular nucleotides to membrane-permeable nucleosides. The encoded protein is used as a determinant of lymphocyte differentiation. Defects in this gene can lead to the calcification of joints and arteries. Two transcript variants encoding different isoforms have been found for this gene.