

## Human S100A12 Antibody Pair Set

<b>Catalog No.</b>	E-KAB-0529	<b>Applications</b>	ELISA
<b>Synonyms</b>	CAAF1;CAGC;CGRP;ENRAGE;MRP-6;MRP6;p6;Calgranulin C		

### Kit components & Storage

Title	Specifications	Storage
Human S100A12 Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze/thaw cycles.
Human S100A12 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze/thaw cycles.

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Product Information

Items		Characteristic (E-KAB-0529)	
		Human S100A12 Capture Antibody	Human S100A12 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human S100A12 protien	Recombinant Human S100A12 protien
	Swissprot	P80511	
Product details	Reactivity	Human	Human
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human S100A12 in ELISAs.	

### For Research Use Only

## Applications

### Human S100A12 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Human S100A12 Capture Antibody	
ELISA Detection	1:1000-1:10000	Human S100A12 Detection Antibody (Biotin)	

**Note:** This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## Background

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein is proposed to be involved in specific calcium-dependent signal transduction pathways and its regulatory effect on cytoskeletal components may modulate various neutrophil activities.

## For Research Use Only