

Recombinant Human MTSS1 Protein (aa1-250, His &MBP Tag)

Catalog Number: PKSH030739

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human MTSS1 protein Met 1-Ser 250, with an N-terminal His & MBP
Calculated MW	71.8 kDa
Observed MW	66 kDa
Accession	EAW92073.1
Bio-activity	Not validated for activity

Properties

Purity	> 80 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 80 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

MTSS1 (Metastasis suppressor 1), also known as Missing in metastasis (MIM), is a tissue-specific regulator of plasma membrane dynamics. MTSS1 is well described for its function as a metastasis suppressor gene and is expressed in a variety of tissues. MTSS1 might be involved in shaping neuronal membranes in vivo. MTSS1 deforms phosphoinositide-rich membranes through its I-BAR domain and interacts with actin monomers through its WH2 domain. MTSS1/MIM was first identified as a metastasis suppressor missing in metastatic bladder carcinoma cell lines. MTSS1 is a prognostic indicator of disease-free survival in breast cancer patients and demonstrates the ability to play a role in governing the metastatic nature of breast cancer cells. MTSS1 may serve as a useful biomarker for the prediction of outcome of gastric cancer. The down-regulation of MTSS1 that may be caused by DNA methylation was also observed in many other types of cancer. Recent work proposed that MIM also potentiates Sonic hedgehog (Shh)-induced gene expression. MTSS1 as a multiple functional molecular player and has an important role in development, carcinogenesis and metastasis.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017