

## Elab Fluor® 488 Anti-Mouse/Rat CD29 Antibody[HMβ1-1]

Catalog Number: E-AB-F1309L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Mouse;Rat
<b>Host</b>	Armenian Hamster
<b>Isotype</b>	Armenian Hamster IgG
<b>Clone No.</b>	HMβ1-1
<b>Isotype Control</b>	[Product E-AB-F09852L]
<b>Conjugation</b>	Elab Fluor® 488
<b>Conjugation Information</b>	Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications


### Recommended usage

**FCM** Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



C57BL/6 murine splenocytes are stained with Elab Fluor® 488 Anti-Mouse/Rat CD29 Antibody (filled gray histogram) or Elab Fluor® 488 Armenian Hamster IgG Isotype Control (empty black histogram).



Rat splenocytes are stained with Elab Fluor® 488 Anti-Mouse/Rat CD29 Antibody (filled gray histogram) or Elab Fluor® 488 Armenian Hamster IgG Isotype Control (empty black histogram).

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	GP1IIaITGB1VLA-β chainβ1 integrin;integrin β1
<b>Uniprot ID</b>	P09055; P49134
<b>Gene ID</b>	16412;24511

### For Research Use Only

## Background

CD29 is a 130 kD protein, also known as integrin  $\beta$ 1, VLA- $\beta$  chain, or GPIIa. It is a member of the integrin family, expressed broadly on leukocytes, endothelial cells, smooth muscle, and epithelial cells. In association with CD49a-f, CD29 forms the VLA-1 through VLA-6 complexes, respectively. It plays an important role in cell-cell or cell-matrix interaction. The HM $\beta$ 1-1 antibody reacts with both mouse and rat CD29. It is able to block cell adhesion and inhibit T cell proliferation.