## **Elabscience**®


## Elab Fluor<sup>®</sup> 488 Anti-Mouse/Rat CD29 Antibody[HMβ1-1]

Catalog Number: E-AB-F1309L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Mouse;Rat
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone No.	ΗΜβ1-1
Isotype Control	[Product E-AB-F09852L]
Conjugation	Elab Fluor <sup>®</sup> 488
Conjugation Information	Elab Fluor <sup>®</sup> 488 is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 $\mu$ L of antibody per test (million cells in 100 $\mu$ L staining volume or per 100 $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



C57BL/6 murine splenocytes are stained with Elab Fluor® 488 Anti-Mouse/Rat CD29 Antibody (filled gray histogram) or Mouse/Rat CD29 Antibody (filled gray histogram) or Elab

Elab Fluor<sup>®</sup> 488 Armenian Hamster IgG Isotype Control (empty black histogram).



Rat splenocytes are stained with Elab Fluor® 488 Anti-

Fluor<sup>®</sup> 488 Armenian Hamster IgG Isotype Control (empty black histogram).

Preparation & Storag	ye
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	GPIIaITGB1VLA-β chainβ1 integrin;integrin β1
Uniprot ID	P09055; P49134
Gene ID	16412;24511

## For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

## **Elabscience**®

Background

CD29 is a 130 kD protein, also known as integrin  $\beta$ 1, VLA- $\beta$  chain, or GPIIa. It is a member of the integrin family, expressed broadly on leukocytes, endothelial cells, smooth muscle, and epithelial cells. In association with CD49a-f, CD29 forms the VLA-1 through VLA-6 complexes, respectively. It plays an important role in cell-cell or cell-matrix interaction. The HM $\beta$ 1-1 antibody reacts with both mouse and rat CD29. It is able to block cell adhesion and inhibit T cell proliferation.