Recombinant Mouse Interleukin-4/IL-4 Protein (His Tag)

Catalog Number: PKSM041094



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Species	Mouse
Mol_Mass	14.6 kDa
Accession	P07750
Bio-activity	Measured in a cell proliferation assay using M-NFS-60 mouse lymphoblast cells. The
	ED ₅₀ for this effect is 0.035 ng/ml.
Duanautias	

	ED ₅₀ for this effect is 0.035 ng/ml.
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.

Please refer to the printed manual for detailed information.

Data

Reconstitution



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-4 (IL-4) is a pleiotropic cytokine that regulates diverse T and B cell responses including cell proliferation, survival and gene expression. IL-4 is produced by mast cells, T cells, and bone marrow stromal cells. IL-4 regulates the differentiation of naive CD4+ T cells into helper Th2 cells, characterized by their cytokine-secretion profile that includes secretion of IL-4, IL-5, IL-6, IL-10, and IL-13, which favor a humoral immune response. Another dominant function of IL-4 is the regulation of immunoglobulin class switching to the IgGl and IgE isotypes. Excessive IL-4 production by Th2 cells has been associated with elevated IgE production and allergic response.

For Research Use Only