

## Recombinant Mouse NOV/CCN3 Protein(His Tag)

**Catalog Number:** PDMM100169

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	Mammalian-derived Mouse NOV/CCN3 proteins Gln22-Ile354, with an C-terminal His
<b>Calculated MW</b>	36.5 kDa
<b>Observed MW</b>	50 kDa
<b>Accession</b>	Q64299
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Mouse NOV/CCN3 proteins, 2 µg/lane of Recombinant Mouse NOV/CCN3 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 50 KD

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.7

Protein NOV homolog, also known as Nephroblastoma-overexpressed gene protein homolog, NOV, and CCN3, is a putative ligand for integrin receptors, is tightly associated with the extracellular matrix, and mediates diverse cellular functions, including cell adhesion and proliferation. CCN3 has been shown to negatively regulate growth although it promotes migration in a cell type-specific manner. This secreted protein belongs to the CCN family, and its expression was observed in a broad variety of tissues from the early stage of development, and altered expression of CCN3 has been observed in a variety of tumors, including hepatocellular carcinomas, Wilm's tumors, Ewing's sarcomas, gliomas, rhabdomyosarcomas, and adrenocortical carcinomas. Mature CCN3 protein has five distinct modules and truncated protein variants with altered function are found in many cancers. CCN3 acts through the core stem cell signaling pathways including Notch and Bone Morphogenic Protein, connecting CCN3 with the modulation of self-renewal and maturation of some cell lineages including hematopoietic, osteogenic, and chondrogenic. CCN3 may affect the extracellular environment of the niche for hematopoietic stem cells. CCN3 has emerged as a key player in stem cell regulation, hematopoiesis, and a crucial component within the bone marrow microenvironment.