

# NFκB-p65 Polyclonal Antibody

catalog number: E-AB-32233

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


|                     |   |
|---------------------|---|
| <b>Reactivity</b>   | Human;Mouse;Rat   |
| <b>Immunogen</b>    | Synthesized peptide derived from human NFκB-p65 around the non-phosphorylation site of Ser281.              |
| <b>Host</b>         | Rabbit  |
| <b>Isotype</b>      | IgG   |
| <b>Purification</b> | Affinity purification   |
| <b>Conjugation</b>  | Unconjugated  |
| <b>buffer</b>       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol. |

## Applications

## Recommended Dilution


|            |              |
|------------|--------------|
| <b>WB</b>  | 1:500-1:2000 |
| <b>IHC</b> | 1:100-1:300  |

## Data




Western Blot analysis of KB, HeLa, 293T, 3T3 cells using NFκB-p65 Polyclonal Antibody at dilution of 1:2000.

**Observed-MV:60 kDa**  
**Calculated-MV:60 kDa**



Immunohistochemistry of paraffin-embedded Mouse kidney tissue using NFκB-p65 Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Human stomach cancer tissue using NFκB-p65 Polyclonal Antibody at dilution of 1:200.

## Preparation & Storage

|                 |  |
|-----------------|--|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| <b>Shipping</b> | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

## Background

## For Research Use Only

# NFκB-p65 Polyclonal Antibody

catalog number: E-AB-32233



Proteins encoded by the v-Rel viral oncogene and its cellular homolog, c-Rel, are members of a family of transcription factors that include the two subunits of the transcription factor NFκB (p50 and p65) and the Drosophila maternal morphogen, dorsal. Both proteins specifically bind to DNA sequences that are the same or slight variations of the 10 bp κB sequence in the immunoglobulin κ light chain enhancer. This same sequence is also present in a number of other cellular and viral enhancers. The DNA binding activity of NFκB is activated and NFκB is subsequently transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct proteins of the same size have been described, designated p105 and p100. The p105 precursor contains p50 at its N-terminus and a C-terminal region that when expressed as a separate molecule, designated pDI, binds to p50 and regulates its activity.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine  
Tel: 400-999-2100

Email: [techsupport@elabscience.cn](mailto:techsupport@elabscience.cn)

Web: [www.elabscience.cn](http://www.elabscience.cn)

Rev. V1.7