Elabscience®


PE/Cyanine5.5 Anti-Human CD18 Antibody[TS1/18.1.2.11]

Catalog Number: E-AB-F1057I

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
lsotype	Mouse IgG1, κ
Clone No.	TS1/18.1.2.11
Isotype Control	PE/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792I]
Conjugation	PE/Cyanine 5.5
Conjugation Information	PE/Cyanine5.5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 690 nm (e.g., a 690/50 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes were stained with PE/Cyanine5.5 Anti-Human CD18 Antibody (filled gray histogram) or PE/Cyanine5.5 Mouse IgG1, κ Isotype Control (empty black histogram).

Keep as concentrated solution.
This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Ice bag
CD18;Cell surface adhesion glycoproteins LFA-1/CR3/p150+95 subunit beta;
Complement receptor C3 subunit beta;Integrin beta-2;Itgb2
P05107

For Research Use Only

Elabscience®

Gene ID Background 3689

CD18 is a 90-95 kD type I transmembrane protein also known as integrin β 2 subunit, LFA-1 β subunit, and β 2 integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains function in adhesion and signaling in hematopoietic cells.