

## Recombinant Human Carboxypeptidase M/CPM Protein (His Tag)

**Catalog Number: PKSH032174**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Carboxypeptidase M;CPM protein Leu18-His422, with an C-terminal His
<b>Calculated MW</b>	47.3 kDa
<b>Observed MW</b>	55 kDa
<b>Accession</b>	P14384
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Concentration</b>	Subject to label value.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/ gel packs. Upon receipt, store it immediately at < - 20°C.
<b>Formulation</b>	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 1mM ZnCl <sub>2</sub> , pH 7.5.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Carboxypeptidase M (CPM) specifically removes C-terminal basic residues (Arg or Lys) from peptides and proteins. Carboxypeptidase exert roles in the physiological processes of blood coagulation/fibrinolysis, inflammation, food digestion and pro-hormone and neuropeptide processing. CPM is believed to play important roles in the control of peptide hormone and growth factor activity at the cell surface, and in the membrane-localized degradation of extracellular proteins. It is widely distributed in a variety of tissues and cells. CPM is involved in peptide metabolism on both the cell surface and in extracellular fluids. CPM functions not only as a protease but also as a binding partner in cell-surface protein-protein interactions.

### For Research Use Only