

Recombinant Mouse TPO/Thpo protein (His Tag)

Catalog Number: PDEM100272

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	E.coli-derived Mouse TPO protein Ser22–Phe195, with an N-terminal His
Calculated MW	19.0 kDa
Observed MW	22 kDa
Accession	P40226
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Thrombopoietin (TPO) is a glycoprotein hormone which belongs to the EPO/TPO family. It produced by the liver and kidney which regulates the production of platelets. Mature mouse Tpo shares 71% and 81% aa sequence homology with human and rat Tpo, respectively. It is an 80-85 kDa protein that consists of an N-terminal domain with homology to Erythropoietin (Epo) and a C-terminal domain that contains multiple N-linked and O-linked glycosylation sites. TPO stimulates the production and differentiation of megakaryocytes, the bone marrow cells that bud off large numbers of platelets. Lineage-specific cytokine affects the proliferation and maturation of megakaryocytes from their committed progenitor cells. It acts at a late stage of megakaryocyte development. It may be the major physiological regulator of circulating platelets.

For Research Use Only