

Ki67 Polyclonal Antibody

catalog number: AN004340L

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Human Ki67 protein expressed by E.coli
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Buffer	PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4


Applications Recommended Dilution

IHC	1:500-1:1000
------------	--------------


Data




Immunohistochemistry of paraffin-embedded human small intestine using Ki67 Polyclonal Antibody at dilution of 1:1000.




Immunohistochemistry of paraffin-embedded human tonsil using Ki67 Polyclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded mouse testis using Ki67 Polyclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded mouse ovary using Ki67 Polyclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded rat ovary using Ki67 Polyclonal Antibody at dilution of 1:1000.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

For Research Use Only

Background

Required to maintain individual mitotic chromosomes dispersed in the cytoplasm following nuclear envelope disassembly. Associates with the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the chromosome surface. Prevents chromosomes from collapsing into a single chromatin mass by forming a steric and electrostatic charge barrier: the protein has a high net electrical charge and acts as a surfactant, dispersing chromosomes and enabling independent chromosome motility. Binds DNA, with a preference for supercoiled DNA and AT-rich DNA. Does not contribute to the internal structure of mitotic chromosomes. May play a role in chromatin organization. It is however unclear whether it plays a direct role in chromatin organization or whether it is an indirect consequence of its function in maintaining mitotic chromosomes dispersed (Probable).

For Research Use Only