



A Reliable Research Partner in Life Science and Medicine

# **Mouse CTSL Antibody Pair Set**

Catalog No. E-KAB-0584 Applications ELISA

**Synonyms** CTSL1;CATL;MEP

## Kit components & Storage

Title	Specifications	Storage
Mouse CTSL Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Mouse CTSL Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0584)	
		Mouse CTSL Capture Antibody	Mouse CTSL Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Mouse CTSL protien	Recombinant Mouse CTSL protien
Information	Swissprot	P06797	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Mouse CTSL in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

### **Applications**

Mouse CTSL Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Mouse CTSL Capture	
Capture		Antibody	10
			Optical Density
ELISA	1:1000-1:10000	Mouse CTSL Detection	Optics
Detection		Antibody (Biotin)	0.1
			100 1000 10000 Mouse CTSL Concentration ( pg/mL )

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## **Background**

Thiol protease important for the overall degradation of proteins in lysosomes (Probable) . Involved in the solubilization of cross-linked TG/thyroglobulin and in the subsequent release of thyroid hormone thyroxine (T4) by limited proteolysis of TG/thyroglobulin in the thyroid follicle lumen. In neuroendocrine chromaffin cells secretory vesicles , catalyzes the prohormone proenkephalin processing to the active enkephalin peptide neurotransmitter. In thymus , regulates CD4+ T cell positive selection by generating the major histocompatibility complex class II (MHCII) bound peptide ligands presented by cortical thymic epithelial cells .

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086
Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a>
Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>