

A Reliable Research Partner in Life Science and Medicine

# PE Anti-Mouse CD161/NK1.1 Antibody[PK136]

Catalog Number: E-AB-F0987UD

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

Reactivity Mouse
Host Mouse

**Isotype** Mouse IgG2a, κ

Clone No. PK136

**Isotype Control** PE Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09803D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

#### Applications Recommended usage


**FCM** Each lot of this antibody is quality control tested by flow cytometric analysis. Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>6</sup> cells is 4.00  $\mu$ L values 1

in 100 µL volume].

# **Data**



C57BL/6 murine splenocytes are stained with PE Anti-Mouse CD161/NK1.1 Antibody and APC Anti-Mouse CD49b Antibody (Left). Unstained splenocytes are used as control.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names CD161 antigen-like family member C;CD161;NK1.1;CD161c;Killer cell lectin-like

receptor subfamily B member 1C;KIrb1c;Ly-55c;NKR-P1 40;NKR-P1.9;NKR-P1C

**Uniprot ID** P27814;P27812;Q99JB4

**Gene ID** 17059;80782

# For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Background**

NK-1.1 surface antigen, also known as CD161b/CD161c and Ly-55, is encoded by the NKR-P1B/NKR-P1C gene. It is expressed on NK cells and NK-T cells in some mouse strains, including C57BL/6, FVB/N, and NZB, but not AKR, BALB/c, CBA/J, C3H, DBA/1, DBA/2, NOD, SJL, and 129. Expression of NKR-P1C antigen has been correlated with lysis of tumor cells in vitro and rejection of bone marrow allografts in vivo. NK-1.1 has also been shown to play a role in NK cell activation, IFN-γ production, and cytotoxic granule release. NK-1.1 and DX5 are commonly used as mouse NK cell markers.

Fax: 1-832-243-6017