Elabscience®


Elab Fluor[®] 700 Anti-Human CD134/OX40 Antibody[Ber-ACT35]

Catalog Number: E-AB-F1153M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	Ber-ACT35
Isotype Control	Elab Fluor [®] 700 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792M1]
Conjugation	Elab Fluor [®] 700
Conjugation Information	Elab Fluor [®] 700 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.





Staining of the 293T cells transfected with pcDNA3.1 plasmid

encoding Human CD134/OX40 gene with Elab Fluor[®] 700 Anti-Human CD134/OX40 Antibody[Ber-ACT35](filled gray

histogram) or Elab Fluor[®] 700 Mouse IgG1, κ Isotype Control(empty black histogram). Total viable cells were used for analysis.

Preparation & Storag	je
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	ACT35 antigen;CD134;OX40L receptor;TNFRSF4;TXGP1L;OX40

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u> Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

Uniprot ID	
Gene ID	
Background	

P43489

7293

CD134, also known as OX40 and TNFRSF4, is a 50 kD type I transmembrane glycoprotein. It is a member of the TNF receptor family. OX40 is expressed on activated T lymphocytes including Th1, Th2, Th17, and Treg cells. The interaction of OX40 with OX40L results in B cell proliferation and antibody secretion, regulation of primary T cell expansion, and T cell survival. OX40 influences the size of the T cell memory pool and regulation of CD4+ T cell tolerance.