A Reliable Research Partner in Life Science and Medicine

EWSR1 Polyclonal Antibody

catalog number: E-AB-52908

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen Fusion protein of human EWSR1

Host Rabbit Isotype IgG


Purification Antigen affinity purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

1:1000-1:5000 WB 1:50-1:300 IHC

Data




EWSR1 Polyclonal Antibody at dilution of 1:1000

Western blot analysis of K562 and HepG2 cell lysates using Immunohistochemistry of paraffin-embedded Human breast cancer tissue using EWSR1 Polyclonal Antibody at dilution of 1:60(×200)

Observed-MW:Refer to figures

Calculated-MW:68 kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using EWSR1 Polyclonal Antibody at dilution of 1:60(×200)

Preparation & Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Storage

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

This gene encodes a multifunctional protein that is involved in various cellular processes, including gene expression, cell signaling, and RNA processing and transport. The protein includes an N-terminal transcriptional activation domain and a C-terminal RNA-binding domain. Chromosomal translocations between this gene and various genes encoding transcription factors result in the production of chimeric proteins that are involved in tumorigenesis. These chimeric proteins usually consist of the N-terminal transcriptional activation domain of this protein fused to the C-terminal DNA-binding domain of the transcription factor protein. Mutations in this gene, specifically a t(11;22)(q24;q12) translocation, are known to cause Ewing sarcoma as well as neuroectodermal and various other tumors. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1 and 14.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Rev. V1.7