A Reliable Research Partner in Life Science and Medicine

# Recombinant Human CD2 Protein (His Tag)

Catalog Number: PKSH033387

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Species Human

**Source** HEK293 Cells-derived Human CD2 protein Lys25-Asp209, with an C-terminal His

Calculated MW22.3 kDaObserved MW34 kDaAccessionP06729

**Bio-activity** Not validated for activity

#### **Properties**

Purity > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.
Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## **Background**

T-cell surface antigen CD2 is also known as Erythrocyte receptor, LFA-2, LFA-3 receptor, Rosette receptor, T-cell surface antigen T11/Leu-5. It is a protein that in humans is encoded by the CD2 gene. It is a single-pass type I membrane protein. contains 1 Ig-like C2-type domain and 1 Ig-like V-type domain. T-cell surface antigen CD2 interacts with lymphocyte function-associated antigen (LFA-3) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. It is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function.