

## Recombinant Human FcRn & B2M Heterodimer (C-6His-Avi) Biotinylated

**Catalog Number:** PKSH033906

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human FcRn;B2M protein Ala24-Ser297&Ile21-Met119, with an C-terminal His & Avi
<b>Calculated MW</b>	31.1&13.5 kDa
<b>Observed MW</b>	35&13 kDa
<b>Accession</b>	AAF72596&P61769
<b>Bio-activity</b>	Loaded Biotinylated Human FcRn Heterodimer (C-6His-Avi) (PKSH033906) on HIS1K Biosensor, can bind Recombinant Human IgG1 Fc (PKSH032558) with an affinity constant of 0.254 uM as determined in BLI assay.


### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 6% Sucrose, 4% Mannitol, 0.05% Tween 80, pH 7.8. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.




Loaded Biotinylated Human FcRn Heterodimer (C-6His-Avi) (PKSH033906) on HIS1K Biosensor, can bind Recombinant Human IgG1 Fc (PKSH032558) with an affinity constant of 0.254 uM as determined in BLI assay.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017



Loaded Biotinylated Human FcRn Heterodimer (C-6His-Avi) (PKSH033906) on HIS1K Biosensor, can bind Recombinant Human IgG2 Fc (223AA) (PKSH033653) with an affinity constant of 0.251  $\mu$ M as determined in BLI assay.

## Background

FcRn complex consist of two subunits: IgGreceptor FcRn large subunit p51 (alpha chain) and Beta-2-microglobulin (Beta chain). The complexes is similar in structure to MHC class I-like heterodimer. Beta-2-microglobulin involved in the presentation of peptide antigens to the immune system. FcRn binds to the Fc region of monomeric immunoglobulins gamma, mediates the uptake of IgG from milk, Possible role in transfer of immunoglobulin G from mother to fetus. A principal component of antibody transport is the neonatal receptor for the Fc portion of immunoglobulin, a heterodimer of a MHC-1 alpha-chain homolog ( FcRn ) and beta-2-microglobulin ( B2M ).