# HASPIN Polyclonal Antibody

**catalog number:** **E-AB-18068**


**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description


| | |
|---|---|
| **Reactivity** | Human |
| **Immunogen** | Synthetic peptide of human HASPIN |
| **Host** | Rabbit |
| **Isotype** | Rabbit IgG |
| **Purification** | Antigen affinity purification |
| **Buffer** | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

| Applications | Recommended Dilution |
|---|---|
| **IHC** | 1:40-1:200 |

## Data



Immunohistochemistry of paraffin-embedded Human tonsil tissue using HASPIN Polyclonal Antibody at dilution of 1:45(×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using HASPIN Polyclonal Antibody at dilution of 1:45(×200)

## Preparation & Storage

| | |
|---|---|
| **Storage** | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| **Shipping** | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

## Background

Serine/threonine-protein kinase that phosphorylates histone H3 at 'Thr-3' (H3T3ph) during mitosis. This positions and activates AURKB and other components of the chromosomal passenger complex (CPC) at centromeres to ensure proper chromatid cohesion, metaphase alignment and normal progression through the cell cycle.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com