

## Recombinant Human CDCP1/CD318 Protein (Fc Tag)

**Catalog Number:** PKSH030671

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                      |   |
|----------------------|---|
| <b>Species</b>       | Human   |
| <b>Source</b>        | HEK293 Cells-derived Human CDCP1/CD318 protein Met 1-Glu343, with an C-terminal mFc |
| <b>Calculated MW</b> | 61.4 kDa  |
| <b>Observed MW</b>   | 89-93 kDa   |
| <b>Accession</b>     | Q9H5V8-3  |
| <b>Bio-activity</b>  | Not validated for activity  |

### Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 90 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from sterile PBS, pH 7.4<br>Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual.             |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

CDCP1 contains three extracellular CUB domains. It is a putative stem cell marker that is highly expressed in some human cancer cells and in both; typical and atypical (cancerous) colons. It interacts with CDH2/N-cadherin; CDH3/P-cadherin; SDC1/syndecan-1; SDC4/syndecan-4 and the serine protease ST14/MT-SP1. It also interacts with SRC and PRKCG/protein kinase C gamma. CDCP1 is taken as a key regulator of EGF/EGFR-induced cell migration. It has been shown that signaling via EGF/EGFR induces migration of ovarian cancer Caov3 and OVCA420 cells with concomitant up-regulation of CDCP1 mRNA and protein. Consistent with a role in cell migration CDCP1 relocates from cell-cell junctions to punctate structures on filopodia after activation of EGFR. It may be involved in cell adhesion and cell matrix association. It also may play a role in the regulation of anchorage versus migration or proliferation versus differentiation via its phosphorylation. It has been taken as a novel marker for leukemia diagnosis and for immature hematopoietic stem cell subsets.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017