

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® Violet 450 Anti-Mouse CD19 Antibody[1D3]

Catalog Number: E-AB-F0986UQ

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Mouse Host Rat

lsotype Rat lgG2a, κ

Clone No. 1D3

Isotype Control Elab Fluor® Violet 450 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833Q]

**Conjugation** Elab Fluor<sup>®</sup> Violet 450

**Conjugation Information** Elab Fluor<sup>®</sup> Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

#### Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>6</sup> cells in 100  $\mu$ L volume].

#### Data



Mouse splenocytes are stained with Elab Fluor<sup>®</sup> Violet 450 Anti-Mouse CD19 Antibody (filled gray histogram). Unstained splenocytes (blank black histogram) are used as control.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names B-lymphocyte antigen CD19;CD19;Cd19;Differentiation antigen CD19

Web: www.elabscience.cn

 Uniprot ID
 P25918

 Gene ID
 12478

### For Research Use Only



## Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

**Background** 

CD19 is a 95 kD glycoprotein also known as B4. It is a member of the lg superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex integral to B cell activation.