Elabscience®


Biotin Anti-Mouse CD31 Antibody[390]

Catalog Number: E-AB-F1180B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
lsotype	Rat IgG2a, κ
Clone No.	390
Isotype Control	Biotin Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833B]
Conjugation	Biotin
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein
	protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow
	cytometric staining, the suggested use of this reagent is $\leq 1.0 \ \mu$ g per 10 ⁶ cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Data



C57BL/6 murine bone marrow cells are stained with FITC Anti-Mouse CD45R/B220 Antibody and Biotin Anti-Mouse

CD31 Antibody followed by Streptavidin-Elab Fluor[®] 647 (Left). Bone marrow cells are stained with FITC Anti-Mouse CD45R/B220 Antibody and Biotin Rat IgG2a, κ Isotype

Control followed by Streptavidin-Elab Fluor[®] 647 (Right).

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	Pecam-1;CD31;PECAM-1;Pecam
Uniprot ID	Q08481
Gene ID	18613

For Research Use Only

Elabscience®

Background

Elabscience Bionovation Inc. A Reliable Research Partner in Life Science and Medicine

CD31 is a 130-140 kD glycoprotein, also known as platelet endothelial cell adhesion molecule (PECAM-1) and EndoCAM. It is a member of the lg superfamily, expressed on endothelial cells, platelets, granulocytes, monocytes/macrophages, dendritic cells, and T and B cell subsets, and is critical for cell-cell interactions. The primary ligands for CD31 have been reported to be CD38 and the vitronectin receptor ($\alpha v \beta 3$ integrin, CD51/CD61). Other reported functions of CD31 are neutrophil emigration to sites of inflammation and angiogenesis.

For Research Use Only