
Recombinant Human ITM2B Protein (His Tag)

Catalog Number: PKSH032599

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human ITM2B protein Tyr76-Ser266, with an C-terminal His
Calculated MW	23.3 kDa
Observed MW	29-33 kDa
Accession	Q9Y287
Bio-activity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel
	packs. Upon receipt, store it immediately at $< -20^{\circ}$ C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Acetate, 10% Trehalose, 25mM NaCl,
	10% Glycerol, 0.05% Tween 80, pH 4.5.
Data	
kDa	MK R
120 90	Augo and a second a



> 95 % as determined by reducing SDS-PAGE.

Background

Integral Membrane Protein 2B (ITM2B) is expressed in the Golgi and on the cell surface. ITM2B forms homodimer through disulfide-linked interaction with SPPL2A, SPPL2B and APP. ITM2B is expressed in brain and the other tissues. Defects in ITM2B cause cerebral amyloid angiopathy ITM2B-related type 1(CAA-ITM2B1) and amyloid angiopathy ITM2B-related type 2(CAA-ITM2B2). CAA-ITM2B1 is characterized by amyloid deposition in the walls of cerebral blood vessels and neurodegeneration in the central nervous system. CAA-ITM2B2 characterized by amyloid deposition in the walls of the blood vessels of the cerebrum, choroid plexus, cerebellum, spinal cord and retina.