A Reliable Research Partner in Life Science and Medicine

# Recombinant Human CCDC47 Protein (His Tag)

Catalog Number: PKSH030669

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Species Human

Source HEK293 Cells-derived Human CCDC47 protein Met 1-Ser135, with an C-terminal His

 Mol\_Mass
 14.7 kDa

 Accession
 Q96A33-1

**Bio-activity** Not validated for activity

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

#### Background

CCDC47 gene is expressed at high level. The gene contains 16 distinct gt-ag introns. Transcription produces 9 different mRNAs, 6 alternatively spliced variants and 3 unspliced forms. There are 3 probable alternative promotors, 3 non overlapping alternative last exons and 8 validated alternative polyadenylation sites. The mRNAs appear to differ by truncation of the 5' end, truncation of the 3' end, presence or absence of a cassette exon, overlapping exons with different boundaries. Functionally, CCDC47 gene has been proposed to participate in processes such as calcium ion homeostasis, embryo development, ER overload response and post-embryonic development. CCDC47 are expected to have molecular function (calcium ion binding) and to localize in various compartments (membrane, endoplasmic reticulu m, integral to membrane, microsome).

#### For Research Use Only

Fax: 1-832-243-6017