

PE Anti-Mouse CD123 Antibody[5B11]

Catalog Number: AN00962D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Mouse
Host	Rat
Isotype	Rat IgG2a
Clone No.	5B11
Isotype Control	PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832D]
Conjugation	PE
Conjugation Information	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications

FCM

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine bone marrow cells with APC Anti-Mouse/Human CD11b Antibody[M1/70] and PE Anti-Mouse CD123 Antibody[5B11] (left) or PE Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	IL-3 Receptor α chain;IL-3Ra
Uniprot ID	Q8CII2

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

Gene ID

16188

Background

CD123 is a 70 kD α chain subunit of the IL-3 receptor (IL-3R α). It is a member of the immunoglobulin superfamily that is expressed on hematopoietic progenitors, basophils, mast cells, and megakaryocytes. This transmembrane glycoprotein can bind IL-3 with low affinity but cannot transduce signals without association with additional protein partners. CD123 can complex with either the common β chain (CDw131) or the IL-3R β chain (AIC2A) to form high-affinity heterodimeric IL-3 receptors. CDw131 can complex with the α subunits of the mouse IL-3R, IL-5R and GM-CSFR to form high-affinity receptors, while the IL-3 R β subunit is specific for IL-3 but binds with low affinity. IL-3 binding to the receptor complex can induce proliferation and differentiation of hematopoietic cells. The 5B11 antibody does not block binding of IL-3 to the high affinity IL-3 receptor.