

Recombinant Human CD16a/FCGR3A Protein (176 Val, His Tag)

Catalog Number: PKSH030288

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	CHO Cells-derived Human CD16a/FCGR3A(176 Val) protein Met 1-Gln 208, with an C-terminal His
Calculated MW	23.3 kDa
Observed MW	40-45 kDa
Accession	AAH17865.1
Bio-activity	Immobilized human CD16a-His at 10 µg/ml (100 µl/well) can bind recombinant human IgG1 (Fc). The EC ₅₀ of human IgG1 (Fc) is 0.40-0.92 µg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.2 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Loaded Recombinant Human CD16a/FCGR3A/Fc gamma RIIIA Protein, His Tag (Cat: PKSH030288) on His1K Biosensor, can bind Rituximab with an affinity constant of 0.46 µM as determined in BLI assay (Sartorius Octet RED384) .

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

The Fc receptor with low affinity for IgG (FCGR3, or CD16) is encoded by 2 nearly identical genes, FCGR3A and FCGR3B, resulting in tissue-specific expression of alternative membrane-anchored isoforms. FCGR3A, it is also known as CD16a, encodes a transmembrane protein expressed on activated monocytes/macrophages, natural killer (NK) cells, and a subset of T cells.

CD16a / FCGR3A is a receptor expressed on NK cells that facilitates antibody dependent cellular cytotoxicity (ADCC) by binding to the Fc portion of various antibodies. CD16a / FCGR3A also has a broader function. CD16a / FCGR3A is directly involved in the lysis of some virus-infected cells and tumor cells by NK cells, independent of antibody binding. Cross-linking of CD16a / FCGR3A on NK cells resulted in increased intracellular Ca²⁺ levels and a cascade of biochemical events similar to those activated by the T cell receptor. CD16a / FCGR3A on human NK cells is a lysis receptor that mediates the direct killing of some virus infected and tumor cells, independent of antibody ligation.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017