



A Reliable Research Partner in Life Science and Medicine

# **Human S100A8 Antibody Pair Set**

Catalog No. E-KAB-0519 Applications ELISA

**Synonyms** 60B8AG;CAGA;CFAG;CGLA;CP-10;L1Ag;MA387;MIF;MRP8;NIF;P8;Calgranulin A

## Kit components & Storage

Title	Specifications	Storage
Human S100A8 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze/thaw cycles.
Human S100A8 Detection Antibody	1 vial, 50 μL	Store at -20℃ for one year.
(Biotin)		Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0519)		
		Human S100A8 Capture Antibody	Human S100A8 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human S100A8 protien	Recombinant Human S100A8 protien	
Information	Swissprot	P05109		
Product details	Reactivity	Human	Human	
	Host	Sheep	Rabbit	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Antigen Affinity	Antigen Affinity	
	Specificity	Detects Human S100A8 in ELISAs.		

For Research Use Only

Tel: 400-999-2100 Web: <a href="www.elabscience.cn">www.elabscience.cn</a> Email: <a href="mailto:techsupport@elabscience.cn">techsupport@elabscience.cn</a>





A Reliable Research Partner in Life Science and Medicine

## **Applications**

Human S100A8 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human S100A8 Capture	
Capture		Antibody	10 M
ELISA Detection	1:1000-1:10000	Human S100A8 Detection Antibody (Biotin)	0.1 0.1 1 10 100  Human S100A8 Concentration (ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calciumbinding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in the inhibition of casein kinase and as a cytokine. Altered expression of this protein is associated with the disease cystic fibrosis. Multiple transcript variants encoding different isoforms have been found for this gene.

Tel: 400-999-2100 Web: <a href="www.elabscience.cn">www.elabscience.cn</a> Email: <a href="techsupport@elabscience.cn">techsupport@elabscience.cn</a>