

(KO Validated) MAP2K2 Polyclonal Antibody

catalog number: E-AB-65103

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide of human MAP2K2 (NP_109587.1).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications


WB	1:500-1:2000
-----------	--------------

Data



Western blot analysis of extracts of HeLa cells using MAP2K2 Polyclonal Antibody at dilution of 1:1000.

Observed-MW:50 kDa
Calculated-MW:44 kDa



Western blot analysis of extracts from normal (control) and MAP2K2 knockout (KO) 293T cells using MAP2K2

Polyclonal Antibody at dilution of 1:1000.

Observed-MW:50 kDa
Calculated-MW:44 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene.

For Research Use Only