

A Reliable Research Partner in Life Science and Medicine

APC Anti-Human IgM Antibody[MHM-88]

Catalog Number: E-AB-F1172E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Mouse Host

Isotype Mouse IgG1, ĸ Clone No. MHM-88

APC Mouse IgG1, k Isotype Control[MOPC-21] [Product E-AB-F09792E] Isotype Control

APC Conjugation

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).


Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with APC Anti-Human IgM Antibody (filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

Preparation & Storage

Keep as concentrated solution. Storage

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names IGHM;Immunoglobulin M;Immunoglobulin heavy constant mu

Uniprot ID P01871 Gene ID 3507

For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Rev. V1.5



Elabscience®

A Reliable Research Partner in Life Science and Medicine

Background

lgM is the first immunoglobulin made by B cells in the immune response. Surface lgM is expressed on immature and mature B cells, while lgM heavy (μ) chain is expressed intracellularly in pre-B cells.