Elabscience®

Purified Anti-Human TCRVy9 Antibody[B3]

catalog number: AN003570P


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Recombinant Human TCRVy9 protein
Host	Mouse
Is otype	Mouse IgG1, ĸ
Clone	B3
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze
	to completely remove the stabilizer prior to labeling.
Applications	Recommended Dilution

FCM

 $2 \,\mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



Human peripheral blood lymphocytes were stained with 0.2 μ g Purified Anti-Human TCRV γ 9 Antibody[B3] (Right) and

 $0.2~\mu g$ Mouse IgG1, κ Isotype Control (Left), followed by

APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles.
Shipping	Ice bag

Background

The V γ 9 TCR is a variant of the TCR γ chain expressed on a subset of γ/δ T cells. V γ 9V δ 2 T lymphocytes, a major γ/δ T cell subset in humans, recognize phosphoantigens, certain tumor cells, and cells treated with aminobisphosphonates. This cell population displays cytolytic activity against various tumor cells. The γ/δ TCR is a heterodimeric TCR complex composed of covalently bound γ and δ chains involved in antigen recognition and the non-covalently associated monomorphic proteins CD3 δ , γ , ε , and ζ chains. The B3 antibody reacts specifically with human TCR V γ 9 as designated by the Lefranc/Foster nomenclature system. Human TCR V γ 9 is also known as TCR V γ 2 under the Strauss, Quertermous nomenclature system. Several TCR γ and δ chain nomenclature systems exist. In order to consolidate the various nomenclature systems, we have created charts for corresponding names across some of the most popular naming methods