MPG Polyclonal Antibody

catalog number: **E-AB-14767**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human
Immunogen	Recombinant protein of human MPG
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
WB	1:1000-1:5000

Data



Western Blot analysis of 231, 293T, Lovo, hepG2, A549, PC3 and A172 cell using MPG Polyclonal Antibody at dilution of 1:1500

Calculated-MW:32 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

Maintenance of DNA sequences is necessary for vertebrates and other life.DNA is under constant stress by a plethora of DNA-damaging agents present in both the environment and within cells.The potentially deleterious effects of DNA lesions in cells are elegantly resolved by sophisticated DNA repair systems, including base excision repair (BER), nucleotide excision repair (NER) and DNA repair methyltransferase (MTase).Methylated bases, such as 3-methyladenine (3MeA) and 7-methylguanine (7MeG) can be formed by agents in the environment and by endogenous cellular processes.Consequently, in the absence of exposure to environmental agents, DNA methylation damage can be incurred on the genomic DNA of normal mammalian cells.DNA N-glycosylases are base excision-repair proteins that locate and cleave damaged bases from DNA as the first step in restoring the sequence.