

## Biotin Anti-Mouse CD4 Antibody[GK1.5]

Catalog Number: E-AB-F1097B

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2b, $\kappa$
<b>Clone No.</b>	GK1.5
<b>Isotype Control</b>	Biotin Rat IgG2b, $\kappa$ Isotype Control[LTF-2] [Product E-AB-F09843B]
<b>Conjugation</b>	Biotin
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

### Applications

#### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per $10^6$ cells in 100 $\mu\text{L}$ volume or 100 $\mu\text{L}$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
------------	---

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	L3T4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;T4
<b>Uniprot ID</b>	P06332
<b>Gene ID</b>	12504
<b>Background</b>	CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, lck.