Elabscience®


APC Anti-Human CD31 Antibody[158-2B3]

Catalog Number: E-AB-F1050E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human	
Host	Mouse	
lsotype	Mouse IgG1, ĸ	
Clone No.	158-2B3	
Isotype Control	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]	
Conjugation	APC	
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an a_{1}	
Storage Buffer	optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter). Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	
Applications	Recommended usage	
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.	

Data



Staining of normal human peripheral blood cells with APC Anti-Human CD31 Antibody[158-2B3] (filled gray histogram) or APC Mouse IgG1, κ Isotype Control (empty black histogram). Cells in the granulocytes gate were used for analysis.

Preparation & Storage	e
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	CD31;EndoCAM;GPIIA';PECA1;PECAM-1;PECAM1;Platelet endothelial cell adhesion molecule

For Research Use Only

Elabscience®

Uniprot ID	P16284
Gene ID	5175
Background	CD31 is a 130-140 kD type I transmendothelial cell adhesion molecule

CD31 is a 130-140 kD type I transmembrane glycoprotein also known as platelet endothelial cell adhesion molecule-1 (PECAM-1) or Endocam. It is expressed on monocytes, platelets, granulocytes, endothelial cells and lymphocyte subsets. CD31 has been reported to bind CD38 and be involved in wound healing, angiogenesis, and cellular migration in an inflammatory situation.

For Research Use Only